

Women and Ischemia Syndrome Evaluation (WISE) Diagnosis and Pathophysiology of Ischemic Heart Disease Workshop

October 2-4, 2002

Session 5

1. Topic and Author

Gender Related Risk Factors for IHD: Novel Risk Factors

Russ Tracy

2. Where we stand in 2002. Overview/rationale for inclusion of topic.

A number of new (novel?) risk factors for IHD have been introduced over the past 10 years¹. These have ranged from new forms of lipid particles, to metabolites, to hemostatic factors, to inflammation markers. Some of these have stood the test of multiple positive studies, while others have not. In many cases they are related and in some cases tightly related. Also, they are commonly related to the set of abnormalities we call the Metabolic Syndrome. This inter-relatedness causes two problems: a) it is often unclear which marker might best represent the group; and b) it is difficult to identify the independent effects of each marker.

Blood proteins associated with inflammation have proven to be consistent risk factors for heart disease in both men and women². The most-studied markers in this class are C-reactive protein, fibrinogen and IL6. These markers are largely independent of lipid effects³, but are closely related to insulin resistance⁴ and adiposity⁵; fibrinogen in particular reflects age⁶ and other measures of existing heart disease⁷. They are also related to coagulation activity, IL-6 and CRP more than fibrinogen. To a certain extent they reflect underlying atherosclerotic disease (fibrinogen⁷ more than CRP⁸), but this relatively weak association⁹ wouldn't appear to account for all, or even most, of their predictive power. While these markers may add independent effects in prediction models which have been fully adjusted for other CVD risk factors, the incremental predictive power is small. However, they can summarize much of the predictive in some other "traditional" risk factors, so that two component risk models (e.g., lipids and CRP³) are quite powerful in prediction.

These associations with heart disease appear to be true for across the life span for both men and women, although predominantly for men in older age groups¹⁰. Some medications have relatively large effects, which may affect the relationship of the marker to disease. For example, post-menopausal estrogen can increase CRP levels 2-3 fold¹¹. The relationship of this effect to actual risk remains unclear. Interestingly, fibrinogen is unaffected by estrogen, or even lowered a bit¹², highlighting that these markers are not all equivalent in all dimensions. It is likely that these markers both reflect underlying pathological processes, and contribute to them as well¹³.

3. Current challenges and the most important issues for future research

What is the best marker, and can it (they) be effectively used in risk prediction?

What are these markers telling us about the underlying pathophysiology and new targets for intervention or prevention?

CVD appears different in some ways in older men and older women; what is the underlying issue(s)?

4. Current challenges in the areas of communicating messages to health care community, patients and the public

The systems we describe as "novel" risk markers are in fact our own defense mechanisms against the "world". For example, blood clotting – it may cause a fatal MI, but we can't live without it. This a complex web of relationships posing real educational challenges.

5. Translating new findings to improved diagnosis and treatment/saving lives.

the obvious target is risk prediction; also, medication use may be helped by measuring these markers.

6. References.

1. Ridker PM, Stampfer MJ, Rifai N. Novel risk factors for systemic atherosclerosis: a comparison of C-reactive protein, fibrinogen, homocysteine, lipoprotein(a), and standard cholesterol screening as predictors of peripheral arterial disease. *Jama*. 2001;285:2481-5.
2. Tracy RP. Inflammation markers and coronary heart disease. *Curr Opin Lipidol*. 1999;10:435-41.

3. Ridker PM. High-sensitivity C-reactive protein: potential adjunct for global risk assessment in the primary prevention of cardiovascular disease. *Circulation*. 2001;103:1813-8.
4. Festa A, D'Agostino R, Jr., Howard G, Mykkanen L, Tracy RP, Haffner SM. Chronic Subclinical Inflammation as Part of the Insulin Resistance Syndrome : The Insulin Resistance Atherosclerosis Study (IRAS). *Circulation*. 2000;102:42-47.
5. Tracy RP. Is visceral adiposity the "enemy within"? *Arterioscler Thromb Vasc Biol*. 2001;21:881-3.
6. Tracy R, Bovill E, Fried L, Heiss G, Lee M, Polak J, Psaty B, Savage P. The distribution of coagulation factors VII, VIII and fibrinogen in adults over the age of 65 years: results from the Cardiovascular Health Study. *Ann. Epidemiol.* 1992;2:509-519.
7. Tracy R, Bovill E, Yanez D, Psaty B, Fried L, Heiss G, Lee M, Polak J, Savage P, for the CHS Investigators. Fibrinogen and factor VIII, but not factor VII, are associated with measures of subclinical cardiovascular disease in the elderly: results from the Cardiovascular Health Study. *Arterioscler. Thromb. Vasc. Biol.* 1995;15:1269-1279.
8. Tracy R, Macy E, Bovill E, Cushman M, Psaty B, Cornell E, Kuller L. Lifetime smoking exposure affects the association of C-reactive protein with cardiovascular disease risk factors and subclinical disease in healthy elderly subjects. *Arterioscler. Thromb. Vasc. Biol.* 1997;17:2167-2176.
9. Folsom AR, Pankow JS, Tracy RP, Arnett DK, Peacock JM, Hong Y, Djousse L, Eckfeldt JH. Association of C-reactive protein with markers of prevalent atherosclerotic disease. *Am J Cardiol*. 2001;88:112-7.
10. Cushman M, Arnold A, Kuller L, Psaty B, Manolio T, Burke G, Polak J, Tracy R. C-reactive protein and risks of myocardial infarction, stroke and death in an elderly cohort. (abstract). *Circulation*. 1999;98:1108.
11. Cushman M, Legault C, Barrett-Connor E, Stefanick ML, Kessler C, Judd HL, Sakkinen PA, Tracy RP. Effect of postmenopausal hormones on inflammation-sensitive proteins: the Postmenopausal Estrogen/Progestin Interventions (PEPI) Study. *Circulation*. 1999;100:717-22.
12. The Writing Group for the PEPI Trial. Effects of estrogen/progestin regimens on heart disease risk factors in postmenopausal women: The Postmenopausal Estrogen/Progestin Interventions (PEPI) Trial. *JAMA*. 1995;273:199-208.
13. Tracy RP. Inflammation in cardiovascular disease: cart, horse, or both? *Circulation*. 1998;97:2000-2.